



**Faculty of Resource Science and Technology**

**PROPAGATION AND ESTABLISHMENT OF *Cassytha filiformis* L.**

**CHER CAMMILLY ANAK ALENG**

**Bachelor of Science with Honours  
(Plant Resource Science and Management)  
2015**

UNIVERSITI MALAYSIA SARAWAK

Grade: \_\_\_\_\_

Please tick (✓)

Final Year Project Report

☒

Masters

☐

PhD

☐

DECLARATION OF ORIGINAL WORK

This declaration is made on the 3 day of 07 year 2015.

Student's Declaration:

I Cher Cammilly anak Aleng (35714) Faculty of Resource Science and Technology  
(PLEASE INDICATE NAME, MATRIC NO. AND FACULTY) hereby declare that the work entitled,  
Propagation and establishment of Cassytha filiformis L. is my original work. I have  
not copied from any other students' work or from any other sources with the exception where due  
reference or acknowledgement is made explicitly in the text, nor has any part of the work been  
written for me by another person.

3/7/2015

Date submitted

Cher Cammilly ak Aleng (35714)

Name of the student (Matric No.)

Supervisor's Declaration:

I, Prof. Dr. Isq bin Ipor (SUPERVISOR'S NAME), hereby certify that the work  
entitled, Propagation and establishment of Cassytha filiformis L. (TITLE) was prepared by the  
aforementioned or above mentioned student, and was submitted to the "FACULTY" as a \*  
partial/full fulfillment for the conferment of Bachelor of Science with Honors Plant Resource  
(PLEASE INDICATE THE DEGREE TITLE), and the aforementioned work, to the best of my  
knowledge, is the said student's work

Received for examination by: mf

(Name of the supervisor)

Date: 3/7/2015

I declare this Project/Thesis is classified as (Please tick (✓)):

- ☐ **CONFIDENTIAL** (Contains confidential information under the Official Secret Act 1972)\*  
☐ **RESTRICTED** (Contains restricted information as specified by the organisation where research was done)\*  
☒ **OPEN ACCESS**

I declare this Project/Thesis is to be submitted to the Centre for Academic Information Services (CAIS) and uploaded into UNIMAS Institutional Repository (UNIMAS IR) (Please tick (✓)):

- ☒ **YES**  
☐ **NO**

#### Validation of Project/Thesis

I hereby duly affirmed with free consent and willingness declared that this said Project/Thesis shall be placed officially in the Centre for Academic Information Services with the abide interest and rights as follows:

- This Project/Thesis is the sole legal property of Universiti Malaysia Sarawak (UNIMAS).
- The Centre for Academic Information Services has the lawful right to make copies of the Project/Thesis for academic and research purposes only and not for other purposes.
- The Centre for Academic Information Services has the lawful right to digitize the content to be uploaded into Local Content Database.
- The Centre for Academic Information Services has the lawful right to make copies of the Project/Thesis if required for use by other parties for academic purposes or by other Higher Learning Institutes.
- No dispute or any claim shall arise from the student himself / herself neither a third party on this Project/Thesis once it becomes the sole property of UNIMAS.
- This Project/Thesis or any material, data and information related to it shall not be distributed, published or disclosed to any party by the student himself/herself without first obtaining approval from UNIMAS.

Student's signature \_\_\_\_\_  
(Date)

Supervisor's signature: mf 3/1/2005  
(Date)

Current Address:

No. 75, Kampung Gram, KM38, Jalan Puncak Borneo, 93250,  
Kuching, Sarawak.

Notes: \* If the Project/Thesis is **CONFIDENTIAL** or **RESTRICTED**, please attach together as annexure a letter from the organisation with the date of restriction indicated, and the reasons for the confidentiality and restriction.

[The instrument was prepared by The Centre for Academic Information Services]

**PROPAGATION AND ESTABLISHMENT OF *Cassytha filiformis* L.**

**CHER CAMMILLY ANAK ALENG**

**(35714)**

A dissertation submitted in partial fulfillment of the requirement for The Degree of Bachelor of  
Science with Honours

Plant Resource Science and Management

Faculty of Resource Science and Technology  
UNIVERSITI MALAYSIA SARAWAK

2015

## APPROVAL SHEET

Name of Candidate : Cher Cammilly Anak Aleng

Title of Dissertation : Propagation and Establishment of *Cassytha filiformis* L.

---

(Prof. Dr. Isa Ipor)

Supervisor

---

(Dr. Rebicca Edward)

Coordinator

Plant Resource Science and Management Programme

Department of Plant Science and Environmental Ecology

Faculty of Resource Science and Technology

University Malaysia Sarawak

## **DECLARATION**

I declare that no portion of the work referred to in this dissertation has been submitted in support of an application for another degree of qualification of this or any other university or institution of higher learning.

---

(CHER CAMMILLY ANAK ALENG)

Programme of Plant Resource Science and Management

Department of Plant Science and Environmental Ecology

Faculty of Resource Science and Technology

University Malaysia Sarawak

## **ACKNOWLEDGEMENTS**

I would like to thank Prof. Dr. Isa Ipor for their advices, motivations and supports during the accomplishment of this project. I am very grateful to all lecturers for their advices given to me and assistance in various ways.

Thank you also to my fellow friends especially to Awang Shahrill, Gracial, Cynthiana, Fiona, Safwan, Hazwan, and Hadzrin for their encouragement and co-operation. I would also like to thank a Master student namely Aisyah Zahidah who guided me with an endless moral support.

Finally, thank you to all my family members wherever they are. I really appreciate their loves and cares forever.

## TABLE OF CONTENTS

Approval Sheet	ii
Declaration	iii
Acknowledgements	iv
Table of contents	v
List of Abbreviation	vii
List of Tables	viii
List of Figures	ix
List of Plates	x
Abstract / Abstrak	xi
1.0 Introduction	1
1.1 Problem statements	3
1.2 Objectives	4
2.0 Literature Review	5
2.1 <i>Cassytha filiformis</i> L.	5
2.2 Importance of <i>Cassytha filiformis</i>	6
2.3 Parasitic Plant	7
2.4 Propagation	8
2.5 Plant Establishment	9
3.0 Materials and Methods	10
3.1 Study site	10
3.2 Propagation Methods	11
3.2.1 Different Placements of <i>Cassytha filiformis</i>	11
3.2.2 Different Weights of <i>Cassytha filiformis</i>	13
3.2.3 Vegetative propagation: Stem Cutting	15



3.2.3.1 Wrapping <i>Cassytha filiformis</i> around the host plant	15
3.2.3.2 Tied Up <i>Cassytha Filiformis</i> to the Host Plant	15
3.3 Statistical Data Analysis	17
4.0 Results	18
4.1 Biomass Assessment of <i>Cassytha filiformis</i>	18
4.1.1 Establishment for Different Biomass Placement	18
4.1.2 Total coverage of <i>Cassytha filiformis</i> 's establishment for different total biomass	20
4.2 Growth Pattern Measurement of <i>Cassytha filiformis</i> Established for Different Placement.	22
4.2.1 Biomass Assessment of <i>Cassytha filiformis</i>	22
4.2.2 Total coverage of <i>Cassytha filiformis</i> 's Establishment for Different Total Biomass.	24
4.3 Vegetative propagation: Stem Cutting	26
4.3.1 Length of <i>Cassytha filiformis</i> , cm	26
4.3.2 Percentage of Success of vegetative propagation through stem Cutting	27
5.0 Disccussions	28
5.1 Different Weight of <i>Cassytha filiformis</i>	28
5.2. Different Placement of <i>Cassytha filiformis</i>	30
5.3 Vegetative propagation: Stem Cutting	32
6.0 Conclusion and Recommendations	34
7.0 References	35
8.0 Appendixes	37

## **LIST OF ABBREVIATIONS**

STDEV - Standard Deviation

SE – Standard Error

MPOB – Malaysian Palm Oil Board

MPOC – Malaysian Palm Oil Council

<b>LIST OF TABLES</b>	<b>PAGE</b>
<b>Table 1</b> Percentage of Success of vegetative propagation through stem cutting	27
<b>Table 2</b> Fresh biomass accessment of propagation through different amount of biomass.	37
<b>Table 3</b> Growth area accessment of propagation through different amount of biomass.	37
<b>Table 4</b> Fresh biomass accessment of propagation through different positions of propagating.	38
<b>Table 5</b> Growth area accessment of propagation through different positions of propagating.	38
<b>Table 6</b> Propagation Method through Cutting.	39

<b>LIST OF FIGURES</b>	<b>PAGE</b>
<b>Figure 1</b> Total weight of established <i>Cassytha filiformis</i> by placing different total biomass of <i>Cassytha filiformis</i> (g).	19
<b>Figure 2</b> Total coverage of <i>Cassytha filiformis</i> propagated in different amount of biomass (m <sup>2</sup> ).	21
<b>Figure 3</b> Total biomass of <i>Cassytha filiformis</i> for different placement.	23
<b>Figure 4</b> Total coverage of <i>Cassytha filiformis</i> propagated in different placement (m <sup>2</sup> ).	25
<b>Figure 5</b> Establishment for different method of cutting.	26

LIST OF PLATES	PAGE
<b>Plate 1</b> The study site of propagation of <i>Cassytha filiformis</i> .	10
<b>Plate 2</b> Propagation method using the different placements of <i>Cassytha filiformis</i> .	12
<b>Plate 3</b> Propagation method using the different initial weight of <i>Cassytha filiformis</i> .	14
<b>Plate 4</b> Propagation method through stem cutting of <i>Cassytha filiformis</i> .	33

# Propagation and Establishment of *Cassytha filiformis* L.

**Cher Cammilly Anak Aleng**

Plant Resource Science and Management  
Faculty of Resource Science and Technology  
University Malaysia Sarawak

## ABSTRACT

*Cassytha filiformis* L. has a potential as a cover crop in oil palm plantation. This study aims to determine the effective method of vegetative propagation for the establishment of *Cassytha filiformis* and to determine the effect of different placements and different weights for the establishment of *Cassytha filiformis*. There are three methods that have been done throughout the experiment. They were different placements of *Cassytha filiformis* on the host plant (*Mikania micrantha*), two different initial weights (20 g and 50 g) of *Cassytha filiformis* placed within the shoot of the host plant (*Mikania micrantha* H. B. K.) and propagation of *Cassytha filiformis* through cutting method using *Asystasia* sp. as the host plant. The results show that the *Cassytha filiformis* established well when it was placed within the host plant with a higher mean of fresh biomass and growth distribution area. *Cassytha filiformis* was also established well when the initial weight for propagation was 50 g with a higher mean of fresh biomass and growth distribution area. Lastly, different cutting methods for propagation showed that *Cassytha filiformis* can be propagated well by wrapping it around of the host plant with a 72.5 % of success. In the end of this experiment, we concluded that *Cassytha filiformis* can be propagated by using its stem and it is established well in their new introduced place.

Keywords: *Cassytha filiformis*, cover crop, parasitic plant

## ABSTRAK

*Cassytha filiformis* L. mempunyai potensi sebagai tanaman penutup bumi di kawasan pertanian seperti ladang kelapa sawit. Kajian ini bertujuan untuk menentukan kaedah pembiakan vegetatif yang berkesan bagi penumbuhan *Cassytha filiformis* dan untuk menentukan kesan kedudukan yang berbeza dan berat yang berbeza bagi penumbuhan *Cassytha filiformis*. Kaedah-kaedah yang digunakan adalah *Cassytha filiformis* akan diletakkan di kedudukan yang berbeza pada tumbuhan perumah (*Mikania micrantha* H. B. K.), dua berat yang berbeza (20 g dan 50 g) *Cassytha filiformis* menggunakan tumbuhan perumah (*Mikania micrantha*) dan pembiakan menggunakan batang *Cassytha filiformis* pada tumbuhan perumah (*Asystasia* sp.). Hasil kajian menunjukkan bahawa penumbuhan *Cassytha filiformis* lebih baik apabila ia diletakkan di kedudukan bawah tumbuhan perumah dengan purata jisim dan keluasan sebaran yang tinggi. Penumbuhan *Cassytha filiformis* juga lebih baik apabila ia ditanam dengan jisim awal sebanyak 50 g menghasilkan purata jisim selepas tuai dan keluasan sebaran yang tinggi. Akhir sekali adalah kaedah keratin batang. Kajian menunjukkan kaedah lilitan pada tumbuhan perumah adalah lebih Berjaya berbanding dengan hanya mengikat di tepi tumbuhan perumah dengan peratusan 72.5 %. Di akhir kajian ini, kesimpulan telah dibuat bahawa *Cassytha filiformis* boleh ditanam menggunakan pembiakan vegetatifnya dan tumbuh dengan baik di tempat baru.

Kata kunci: *Cassytha filiformis*, parasit, tanaman penutup bumi

## 1.0 INTRODUCTION

Malaysia is one of the biggest producers and exporters of oil palm and oil palm products which make Malaysia have an important role to play in fulfilling the global need for oils and fats sustainably (MPOC, nd). According to MPOC (nd) approximately 4.49 million hectares of land in Malaysia is under oil palm cultivation. MPOB (nd) stated that in 2011, the total planted area was 4.917 million hectares. This shows that palm oil has become the most important crop in Malaysia. The Malaysian oil palm industry can easily meet the local oils and fats demand and the excess kernel oil produced will be exported (MPOB, nd).

As the palm oil plantation getting more important in Malaysia, weeds have become a major problem in the palm oil plantation area. Essandoh et al. (2011) stated that weeds such as *Mikania micrantha*, *Mikania cordata*, *Chromolaena odorata*, *Imperata cylindrical* and *Asystasia* sp. compete with the oil palm for nutrients, moisture and sunlight which can cause yield depression. Sustainable weed management in oil palm plantation has been a challenge now a day (Samedani et al., 2014). According to Ishak et al. (2011), there is possible way to control the weeds problem but the required labor for manual weeding is costly and hardly available and the agricultural workers who performing manual weeding are exposed to high risk of musculoskeletal disorders to the lower back.

Cultivation of cover crops qualifies as part of a sustainable agricultural practice (Samedani et al., 2014). Leguminous cover crops are used to co-exist with the oil palm following jungle clearing and planting or replanting, to provide complete cover to an otherwise bare soil to protect the soil from the forces of erosion. The leguminous cover crops also have multiple functions such as reducing soil water evaporation, runoff losses, soil erosion, improve or maintain soil fertility and recycling of nutrients. The commonly

used leguminous cover crops species in Malaysia are *Pueraria phaseloides* (synonym for *Pueraria javanica*), *Centrosema pubescens*, *Calopogonium mucunoides*, *C. Caeruleum* and *Mucuna bracteata* (Samedani et al., 2014). However, seeds of the cover crop can be expensive and it may not germinate due to inappropriate pre-sowing treatment caused by human error. To overcome this problem, parasitic plant is suggested used as an alternative cover crop.

Study has been done on identifying the propagation and establishment of a parasitic plant specifically *Cassytha filiformis*. This plant depends on its host plant in order to survive. The host plant plays important roles to provide enough water, food and mineral for the parasitic plant to survive. The *Cassytha filiformis* usually attack woody host such as citrus, mango (*Mangifera indica*), cloves (*Eugenia aromatic*), nutmeg (*Myristica fragrans*) and avocado (*Persea Americana*) (Nelson, 2008). In this research, *Cassytha filiformis* were propagate using the vegetative part which is stem and establishment was observed.



## 1.1 Problem Statements

Cultivation of cover crops is one of a sustainable agricultural practice in crop plantation (Samedani et al., 2014). However, seeds of the cover crop can be expensive and it may not germinate due to inappropriate pre-sowing treatment caused by human error. In order to overcome this problem, we suggest using the parasitic plant as an alternative cover crop. *Cassytha filiformis* is a parasitic plant which can be used as a cover crop plant. *Cassytha filiformis* also has a lot of benefit to human being. *Cassytha filiformis* is commonly propagated by using seeds. Therefore, for this study, we propagated the *Cassytha filiformis* by using its stem.

## 1.2 Objectives

1. To determine the effective method of vegetative propagation for the establishment of *Cassytha filiformis*.
2. To determine the effect of different placement and different weights for the establishment of *Cassytha filiformis*.

## **2.0 LITERATURE REVIEW**

### **2.1 Botanical descriptions of *Cassytha filiformis* L.**

*Cassytha filiformis* L. is a leafless, climbing, twining, vine-like, autoparasitic and plant hyperparasitic phanerogam (seed-bearing plant) in the plant family Lauraceae. According to Mythili et al. (2011) *Cassytha filiformis* is a leafless and perennial vine with small scales as a replacement of the leaves and adheres to the host by the haustoria that penetrates the epidermis of the other plants, and hence helps to spread through the tops of the trees and bushes, thus ultimately forming a long festoons of colourful greenish yellow vines. The individual stems that are copiously branched have a range of 1 to 3 millimeters in the diameter and finally attain a maximum length of 10 to 20 feet. The flowers were produced all over the year with a bisexual nature with a stamen and perianth arrangement that looks similar to the ordinary avocado. The ovary is the one first getting exposed and later on becomes enveloped by the enlargement and then the over growth of the calyx tube occurs. The fruit is about the size of the large pea and gets closely enclosed by means of the succulent calyx.

## 2.2 Importance of *Cassytha filiformis*

The *Cassytha filiformis* has an active components in which can be used for medicinal purposes. Mythili et al. (2011) stated that *Cassytha filiformis* has been reported as a beneficial medicine against the gonorrhea, kidney ailments and as the diuretic. In Africa it was used to treat the cancer and African trypanosomiasis. Nelson (2008) also had mention that *Cassytha filiformis* is widely used for its medicinal purposes such as in the Pasific region. They use the plant to treat the jellyfish stings in Fiji. The Hawaiians used this parasitic plant as a ceremonial human ornament and also in the decorative garlands and in the lei. The brown color of the stem is used as the coloring agent (neutral dye) as it provided brown color for a fabric (Mythili, 2011). *Cassytha filiformis* is also a source of food for human and animal. It is also can be a potential biological control agent for invasive plant since it is a parasitic plant (Nelson, 2008). On the other hand, it is also a pestiferous and pathogenic weed which is a major threat to the agriculture and other endangered plants as they are capable of transmitting the pathogens between the plants (Mythili et al., 2011).

*Cassytha filiformis* is a parasitic plant (Nelson, 2008). This means this plant is harmful to its host plant. A disease of the host plant is caused by the effect of infection of host plants by its haustoria which are found on the stem. The Haustoria will penetrate the host epidermis and extend into more interior tissue, extracting cellular nutrients and water from plant phloem and xylem. This will make the *Cassytha filiformis* absorb nutrients and water from the host plant. These activities do not create immediate, fatal damage to the host plant.

## 2.3 Parasitic Plant

Parasite is from a Greek. Para means beside and sitos means grain or food which literally means “beside the food”. The lifecycle of parasitic plant species is closely regulated by the presence of their host plant and signaling molecules released by the host plant play an important role in this interaction (Bouwmeester, 2013). Parasitic plant may cause damage to their host plant which may be one of the demand economic plants.

Most of plant can produce their own food because it contains chlorophyll but it is different for the parasitic plant. The parasitic plant does not has a chlorophyll which cause it cannot produce its own food and depending 100% to its host plant for the food source.

Daniel L. et al. (2010) define a parasitic plant as an angiosperm that directly attaches to another plant via haustorium. Haustorium is a modified root that forms a morphological and physiological link between the parasite and its host plant. A parasitic plant obtains a mature, functioning root system when their seedling forms a haustorium. For the root parasitism, it prefers an open, sunny area. These areas are usually containing dense groundcover vegetation which a great amount of host plant for the parasite plant.

## 2.4 Propagation of *Cassytha filiformis*

During winter, *Cassytha filiformis* seeds germinate and produce a slender yellowish shoot, but not root and when it is in contact with a susceptible host, the shoot will encircle the host plant, haustoria is formed and it begins to climb up the plant (Mukhtar et al., 2010).

Seedlings of *Cassytha filiformis* can survive for up to two months without a host and growing to a length of 30 cm or more. Nelson (2008) stated that *Cassytha* species are parasitic vines with small haustoria. Haustoria is a modified stem which helps the parasitic plant to absorb nutrients and water from the host plant. The haustoria of *Cassytha filiformis* penetrate the host epidermis and extend into more interior tissues, extracting cellular nutrients and water from plant phloem and xylem.

Nelson (2008) also mentioned that two *Cuscuta* species of the family Convolvulaceae have a similar appearance to *Cassytha filiformis* and also have a similar mode of parasitism. *Cuscuta sandwichiana* is being propagated by using seeds same as *Cassytha filiformis* and it can also be grown by wrapping cuttings around host plants such as weeds or other unwanted plants (Hawaiian Native Plant Propagation Database: *Cuscuta sandwichiana*, 2002)

## **2.5 Plant Establishment**

Plant establishment is defined as the process by which a plant becomes established in a new habitat. General environment can be a difficult one for establishing some native plants. Plants need a necessary time to adapt well to their new environment. The new environment may be much extreme growing conditions than their origin environment.

The plant can dry out easily because the root are not deep into the natural soil to consume water and nutrient, there is no sufficient plant mass to handle a bad attack by pest and it need to overcome 'transplant shock' where it may be damage particularly its root in planting process (Plant Establishment Methods, 2011). A new plant has a greater change of things going wrong than one that is well established. Plants will be considered established once they have rooted into the growing medium and produce new growth. Much care needed to be considered during the plant establishment period.

### 3.0 MATERIALS AND METHODS

#### 3.1 Study Site

The *Cassytha filiformis* were taken from Unijaya, UNIMAS. The research was carried out at Tun Ahmad Zaidi College, UNIMAS and Kampung Giam, Padawan.



1(a)



1(b)

**Plate 1.** The study site of propagation of *Cassytha filiformis*. 1(a) source of *Cassytha filiformis*. 1(b) The study site.